

High Energy and Momentum Resolved Photoemission Studies of Quasi-One-Dimensional Blue Bronze $K_{0.3}MoO_3$

Physics Department, BNL:

Alexei Fedorov

Peter Johnson

V.N. Muthukumar

Sergei Brazovskii

Physics Department, Boston University:

Jinyu Xue

Laurent Duda

Kevin Smith

Chemistry Department, Rutgers University:

Martha Greenblatt

William McCarrol

NSLS, BNL:

Steven Hulbert

Supported by the Department of Energy under DE-AC02-98CH10886

**Why are we interested in
high energy and momentum resolution?
What are the goals?**

Nesting properties of the Fermi surfaces
/Charge density waves/

Photoemission spectral functions $A(k,w)$
/direct comparison with theoretical predictions/

Outline

Experimental details:

- ✓ **Photoelectron spectrometer**

Introduction to $K_{0.3}MoO_3$:

- ✓ **Crystal structure**
- ✓ **Electronic structure**
- ✓ **Structural studies /Charge Density Waves/**

Experimental data:



- ✓ **Band structure of $K_{0.3}MoO_3$**
- ✓ **Fermi wave vectors versus temperature**
- ✓ **Incommensurate to commensurate CDW transition**
- ✓ **Signatures of non-Fermi liquid behavior**

Photoelectron Spectrometer

SES-200: 200 millimeters hemispherical deflector capable of multichannel detection in emission angle and kinetic energy

Example of angle resolved data:



*$h\nu = 21.22\text{ eV}/\text{He I radiation}/$
 $\text{Cu}(111)$, bulk bands and sp surface state*



- ✓ *Energy resolution $\sim 10\text{ meV}$*
- ✓ *Angle resolution $\sim 0.2^\circ$*
- ✓ *Base pressure $\sim 2 \times 10^{-11}\text{ Torr}$*

Presently located at the undulator beamline U13UB at the National Synchrotron Light Source

Low dimensionality \Rightarrow { Charge Density Waves (CDW) /Peierls transitions /
Electron correlation effects
/non-Fermi liquid behavior, spin-charge separation, HTSC/





J.-P. Pouget et al., J. Physique Lett. 44, L113 (1973)

Electronic structure of $K_{0.3}MoO_3$

/tight-binding calculations/

M.-H. Whangbo and L.F. Schneemeyer, Inor. Chem. 25, 2424 (1986)



$$|I\rangle = \frac{1}{\sqrt{2}} (|\Phi_a\rangle + |\Phi_b\rangle)$$

$$|II\rangle = \frac{1}{\sqrt{2}} (|\Phi_a\rangle - |\Phi_b\rangle)$$

Two bands crossing the Fermi level
How many Charge Density Waves?


Structural studies of CDW in $K_{0.3}MoO_3$

/Single Charge Density Wave/

(i) Diffuse X-ray scattering

$$/q_{CDW} = 2k_F b^x /$$

J.-P. Pouget et al.





Nesting:

Fermi surface of the first band
is nested to the Fermi surface
of the second band

(ii) Temperature dependent neutron scattering

/incommensurate to commensurate transition/

M.Sato, H. Fujishita and S.Hoshito,
J. Phys. C: Solid State phys., 16, L877 (1983)



CDW wave vector
 $q_{CDW} : k_{F1} + k_{F2}$

Temperature dependence of CDW wave vector:


- ◊ Thermally activated charge transfer between bands crossing the Fermi level and third band above it
/Pouget et al./
- ◊ Shift of the chemical potential
/Pouget & Nougera, Artemenko et al./
- ◊ Hidden temperature dependence of the nesting vector
/Intention of the present study/

Goals of photoemission experiment:

- ◊ Direct monitoring k_{F1} and k_{F2}
- ◊ Temperature dependence of $(k_{F1}+k_{F2})$

Direct monitoring electron bands in $K_{0.3}MoO_3$

/3-D maps of photocurrent/




Experimental details:

Samples cleaved *in situ*

Liquid He cryostat provides temperatures from ~20 K to ~450 K


Temperature monitored with a help of OMEGA CY7 sensor

Momentum Distribution Curves at E_F




Incommensurate to commensurate CDW transition in $K_{0.3}MoO_3$

/comparing neutron scattering data with nesting vector measured in photoemission experiment/




Fermi surface of an array of coupled chains

/tight binding calculation/



Fermi surface is given by:

$$\mu = -2\cos(k_{\parallel}) \pm (t_{\perp} + 2t_{\perp} t \cos(k_{\perp}) + t)^{\frac{1}{2}}$$



What are the signatures of non-Fermi liquid behavior in photoemission?

Spin-charge separation \Rightarrow $\left\{ \begin{array}{l} \text{Observation of two dispersing features} \\ \text{corresponding to the charge and spin} \\ \text{degrees of freedom} \end{array} \right.$

Breakdown of the quasiparticle picture \Rightarrow $\left\{ \begin{array}{l} \text{Suppression of the spectral weight} \\ \text{at the Fermi energy} \end{array} \right.$

(a) *Effect of disorder on
differential conductance*
Hans J. Kwock and Michael H. Szymanski



Order-disorder transitions of the topological insulator

W. MARCHIAFAVA

SIAM JOURNAL ON APPLIED MATHEMATICS

II-661 JOURNAL OF APPLIED MATHEMATICS



APPLIED COMMUNICATIONS



Charge–spin separation and the spectral properties of Luttinger liquids


Johannes Voit

Institut Laue–Langevin, BP 156, 38042 Grenoble Cédex 9, France






Anomalous Scaling and Spin-Charge Separation in Coupled Chains

Peter Kopietz, Volker Meden, and Kurt Schönhammer


Institut für Theoretische Physik der Universität Göttingen, Bunsenstrasse 9, D-37073 Göttingen, Germany
(Received 19 August 1994)

Spectral function in $K_{0.3}MoO_3$ at the Fermi wave vector










Suppression of spectral weight in photoemission from low-dimensional conductors: influence of momentum resolution



Spectral line-shapes versus electron momentum



Spectral line-shapes versus temperature



Gap opening at the
Fermi wave vector

